

## Account of Non-Diagonal $d$ - $d$ -Electron Couplings in Fe-Co Liquid Alloy

Nikolay Dubinin<sup>1,2</sup>

<sup>1</sup>) Ural Federal University, Mira st. 19, 620002 Ekaterinburg, Russia

<sup>2</sup>) Institute of Metallurgy of the Ural Branch of the Russian Academy of Sciences,  
Amundsen st. 101, 620016 Ekaterinburg, Russia  
[ned67@mail.ru](mailto:ned67@mail.ru)

Copyright © 2013 Nikolay Dubinin. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

The partial Wills-Harrison effective pair potentials in liquid equiatomic Fe-Co alloy are considered. It is shown that the account of the non-diagonal  $d$ - $d$  couplings between electrons leads to the same changes in characteristics of the pair-potential first minimum as in the case of the pure transition metals.

**Keywords:** liquid Fe-Co alloy, transition metal, Wills-Harrison model,  $d$ -state coupling


The Wills-Harrison [1] partial effective pair potentials,  $\varphi_{ij\text{WH}}(r)$ , between atoms Fe and Co in equiatomic Fe-Co alloy at  $T=1863\text{K}$  are calculated at different values of the suggested in [2] probability  $p$  that not only diagonal  $d$ - $d$  couplings are possible at condition that all  $d$ - $d$  couplings (diagonal and non-diagonal) are equiprobable in this case.

Following [3] we use the local Bretonnet-Silbert (BS) model pseudopotential [4] extended to binary alloys in [5] for description the  $s$ -electron contribution,  $\varphi_{sij}(r)$ , to  $\varphi_{ij\text{WH}}(r)$ .

As follows from the previous paper,  $\varphi_{ij\text{WH}}(r) = \varphi_{sij}(r)$  at  $p = 1$ . It denotes here that  $\varphi_{ij\text{WH}}(r) = \varphi_{ij\text{BS}}(r)$  at  $p = 1$ .

It is clear from Fig.1 that changing  $p$  from 0 up to 0.5 slightly influences the depth and position of the first minimum of  $\varphi_{ij\text{WH}}(r)$ . At the same time, these characteristics become strong different at  $p = 1$ . Such a tendency is the same as it

was observed in the case of pure transition metals [2].



**Figure 1.**  $\varphi_{ijWH}(r)$  between atoms Fe and Co in liquid equiatomic Fe-Co alloy ( $p = 0$  – solid line;  $p = 0.5$  – dotted line;  $p = 1$  – dashed-dotted line).

## References

- [1] J.M. Wills, W.A. Harrison, Interionic interactions in transition metals, *Phys. Rev. B*, 28 (1983), 4363-4373.
- [2] N.E. Dubinin, Account of non-diagonal coupling between  $d$  electrons at describing the transition-metal pair potentials, *J. Phys.: Conf. Series*, 338 (2012), 012004.
- [3] N.E. Dubinin, L.D. Son, N.A. Vatolin, The Wills-Harrison approach to the thermodynamics of binary liquid transition-metal alloys, *J. Phys.: Condens. Matter*, 20 (2008), 114111.
- [4] J.L. Bretonnet, M. Silbert, Interionic interactions in transition metals. Application to vanadium, *Phys. Chem. Liq.*, 24 (1992), 169-176.
- [5] N.E. Dubinin, L.D. Son, N.A. Vatolin, Thermodynamic properties of liquid binary transition-metal alloys in the Bretonnet-Silbert model, *Defect and Diffusion Forum*, 263 (2007), 105-110.

**Received: April 30, 2013**